

University of Groningen

## Resolution and circular dichroism of an asymmetrically cage- opened [60]fullerene derivative

Hummelen, J.C.; Keshavarz, K.; van Dongen, J.L.J.; Janssen, R.A.J.; Meijer, E.W.; Wudl, F.; Keshavarz-K, Majid

*Published in:*  
Chemical Communications

*DOI:*  
[10.1039/a707274k](https://doi.org/10.1039/a707274k)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
1998

[Link to publication in University of Groningen/UMCG research database](#)

### *Citation for published version (APA):*

Hummelen, J. C., Keshavarz, K., van Dongen, J. L. J., Janssen, R. A. J., Meijer, E. W., Wudl, F., & Keshavarz-K, M. (1998). Resolution and circular dichroism of an asymmetrically cage- opened [60]fullerene derivative. *Chemical Communications*, 1(2), 281 - 282. <https://doi.org/10.1039/a707274k>

### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

# Resolution and circular dichroism of an asymmetrically cage-opened [60]fullerene derivative

Jan C. Hummelen,<sup>\*a</sup> Majid Keshavarz-K,<sup>b</sup> Joost L. J. van Dongen,<sup>c</sup> René A. J. Janssen,<sup>c</sup> E. W. Meijer<sup>c</sup> and Fred Wudl<sup>b</sup>

<sup>a</sup> Department of Organic and Molecular Inorganic Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

<sup>b</sup> Institute for Polymers and Organic Solids, University of California at Santa Barbara, Santa Barbara, CA 93106, USA

<sup>c</sup> Laboratory of Organic Chemistry, Eindhoven University of Technology, 5600 MB Eindhoven, The Netherlands

The enantiomers of the first cagewise *inherently* asymmetric C<sub>60</sub> derivative, *i.e.* N-MEM (MEM = 2-methoxyethoxy-methyl) keto lactam **1**, have been separated by HPLC and their chiroptical properties compared with data obtained from C<sub>60</sub> derivatives that are dissymmetric or asymmetric due to chiral addends.

N-MEM keto lactam **1** (Fig. 1) is the first example of a well-defined open-cage derivative of C<sub>60</sub>.<sup>1</sup> It is obtained from C<sub>60</sub> in two synthetic steps, and (±)-**1** serves as the key intermediate in the synthesis of azafullerenes, *e.g.* (C<sub>59</sub>N)<sub>2</sub>,<sup>2</sup> C<sub>59</sub>NH<sup>3</sup> and K<sub>6</sub>C<sub>59</sub>N.<sup>4</sup> Stereochemically, **1** is an interesting molecule, because it can be regarded as a (highly symmetrical) sphere with a chiral orifice. A larger version could be envisioned as a chiral selector, operating by enantioselective endohedral complexation.

Various chiral C<sub>60</sub> derivatives have been prepared thus far.<sup>5,6</sup> Chirality can be introduced into C<sub>60</sub> derivatives *via* chiral addends, either asymmetric<sup>7,8</sup> or C<sub>2</sub>-symmetric,<sup>9</sup> as well as *via* an inherently asymmetric addition or substitution pattern, such as in 2:1 adducts with C<sub>2</sub>-symmetry.<sup>10,11</sup> Higher fullerenes and carbon nanotubes can be inherently chiral.<sup>12</sup> C<sub>2</sub> C<sub>76</sub>, D<sub>3</sub> C<sub>78</sub> and D<sub>2</sub> C<sub>80</sub> have been obtained in enantiomerically pure form *via* kinetic resolution by asymmetric osmylation, providing optically active forms of carbon.<sup>13</sup> Since keto lactam **1** is the first open-cage C<sub>60</sub> derivative with an inherently asymmetric cage functionalization pattern, it is of interest to determine a possible resemblance of the chiroptical properties of **1** with those of the chirally modified C<sub>60</sub> compounds. Here we present the successful resolution of (±)-**1** and the chiroptical properties of the individual enantiomers.

Chromatography of racemic **1** as a 0.3 mg ml<sup>-1</sup> solution in toluene–1,2-dichlorobenzene (2:1 v/v) on an analytical chiral stationary phase HPLC column [Bakerbond Pirkle Type DNBPG 5 μm (4.6 × 250 mm)], using *n*-hexane–CHCl<sub>3</sub>–Pr<sup>i</sup>OH (70:30:1 v/v/v) as the eluent (flow rate 1 ml min<sup>-1</sup>) and UV detection at λ = 328 nm, gave enough separation to yield the enantiomers in 80 and 92% ee (as inferred from a second HPLC experiment, Fig. 2), respectively. The two fractions were




Fig. 1 Molecular structure of the two enantiomeric forms of **1** with configurational description <sup>1</sup>C and <sup>1</sup>A according to ref. 6

assigned to (+)-**1** and (–)-**1**, respectively, by determining their specific rotations at 589 nm.

The UV–VIS spectrum of (±)-**1**, the circular dichroism (CD) spectrum and the resulting *g* value ( $g = \Delta\epsilon/\epsilon \approx \Delta A/A$ ) of (+)-**1** are shown in Fig. 3. The CD spectrum of (–)-**1** was found to have a mirror image relation to that of (+)-**1**. Similar to the UV–VIS absorption, the CD spectrum of (+)-**1** extends throughout the 200–700 nm spectral range. The strongest Cotton effect is found at λ = 325 nm ( $\Delta\epsilon = +29 \text{ M}^{-1} \text{ cm}^{-1}$ ), coinciding with the lowest energy dipole-allowed transition of **1** (λ = 328 nm). Comparison with, for example, (+)-hexahelicene, which has a *g* value of  $+7.0 \times 10^{-3}$  at λ = 325 nm for the π–π\* transition, shows however that the *g* value of (+)-**1** ( $g_{325} = +6.4 \times 10^{-4}$ ) is an order of magnitude lower and represents a typical value for a noninherently dissymmetric chromophore.<sup>14</sup> Various chirally modified [6,6]-dihydrofullerene derivatives show a diagnostic CD band at about λ = 430 nm, whose sign has been taken as an indicator for the absolute configuration.<sup>8,15–17</sup> For (+)-**1**, a local extreme in the CD spectrum is found at λ = 426 nm, where the Cotton effect is negative. Tentatively applying the sector rule of Wilson *et al.* to **1** suggests that (+)-**1** corresponds to the <sup>1</sup>A configuration (Fig. 1).<sup>15</sup> The largest *g* values of (+)-**1** are found in the wavelength range λ = 600–700 nm, with a maximum of  $g = -1.7 \times 10^{-2}$  at 659 nm, a typical value for a magnetically-allowed dipole-forbidden transition.

In summary, we have separated the two enantiomers of the C<sub>60</sub> derivative (±)-**1** using chiral HPLC, providing a first example of an open-cage fullerene with a chiral orifice. The inherently chiral functionalization of the eleven-membered ring causes a dissymmetric perturbation on the π system of the fullerene, resulting in optical activity extending over the entire





Fig. 2 Chiral HPLC chromatogram of separated samples of (a) (+)-**1** and (b) (–)-**1** (Bakerbond Pirkle Type DNBPG, eluent: *n*-hexane–CHCl<sub>3</sub>–Pr<sup>i</sup>OH (70:30:1 v/v/v), flow rate 1 ml min<sup>-1</sup>, UV detection at λ = 328 nm)



**Fig. 3** (a) UV–VIS absorption spectrum of (±)-**1** ( $c = 7.6 \times 10^{-6}$  M) [ $\lambda_{\text{max}}/\text{nm} = 204$  ( $\epsilon = 1.5 \times 10^5 \text{ M}^{-1} \text{ cm}^{-1}$ , *n*-hexane), 260 ( $1.2 \times 10^5$ ,  $\text{CHCl}_3$ ), 328 ( $4.5 \times 10^4$ ,  $\text{CHCl}_3$ ), 428 ( $4.7 \times 10^3$ ,  $\text{CHCl}_3$ ) and 688 ( $4.2 \times 10^2$ ,  $\text{CHCl}_3$ )]. (b) CD spectra (left axis) and *g* value (right axis) of (+)-**1** ( $c = 7.6 \times 10^{-6}$  M). The UV–VIS and CD spectra were recorded using a solution in *n*-hexane for the region  $\lambda = 200$ –248 nm and in  $\text{CHCl}_3$  for the region  $\lambda = 248$ –800 nm. The  $\text{CHCl}_3$  spectra were recorded at higher concentrations and mathematically scaled to the spectrum in *n*-hexane for representation.

absorption spectrum, with high *g* values in the low-energy region.

Dedicated to Professor Dr Hans Wynberg on the occasion of his 75th birthday on 28th November 1997.

## Footnote and References

\* E-mail: j.c.hummelen@chem.rug.nl

- 1 J. C. Hummelen, M. Prato and F. Wudl, *J. Am. Chem. Soc.*, 1995, **117**, 7003.
- 2 J. C. Hummelen, B. Knight, J. Pavlovich, R. González and F. Wudl, *Science*, 1995, **269**, 1554.
- 3 M. Keshavarz-K., R. González, R. G. Hicks, G. Srdanov, V. I. Srdanov, T. G. Collins, J. C. Hummelen, C. Bellavia-Lund, J. Pavlovich, F. Wudl and K. Holczer, *Nature*, 1996, **383**, 147.
- 4 K. Prassides, M. Keshavarz-K., J. C. Hummelen, W. Andreoni, P. Giannozzi, E. Beer, C. Bellavia, L. Cristofolini, R. González, A. Lappas, Y. Murata, M. Malecky, V. I. Srdanov and F. Wudl, *Science*, 1996, **271**, 1833.
- 5 F. Diederich, C. Thilgen and A. Herrmann, *Nachr. Chem. Tech. Lab.*, 1996, **44**, 9.
- 6 C. Thilgen, A. Herrmann and F. Diederich, *Helv. Chim. Acta*, 1997, **80**, 183.
- 7 A. Vasella, P. Uhlmann, C. A. A. Waldruff, F. Diederich and C. Thilgen, *Angew. Chem.*, 1992, **104**, 1388.
- 8 A. Bianco, M. Maggini, G. Scorrano, C. Toniolo, G. Marconi, C. Villani and M. Prato, *J. Am. Chem. Soc.*, 1996, **118**, 4072.
- 9 M. Maggini, G. Scorrano, A. Bianco, C. Toniolo and M. Prato, *Tetrahedron Lett.*, 1995, **36**, 2845.
- 10 J. M. Hawkins, A. Meyer and M. Nambu, *J. Am. Chem. Soc.*, 1993, **115**, 9844.
- 11 A. Hirsch, I. Lamparth and H. R. Karfunkel, *Angew. Chem.*, 1994, **106**, 453.
- 12 F. Diederich and R. L. Whetten, *Acc. Chem. Res.*, 1992, **25**, 119; S. Iijima, *Nature*, 1991, **354**, 56.
- 13 J. M. Hawkins and A. Meyer, *Science*, 1993, **260**, 1918; J. M. Hawkins, M. Nambu and A. Meyer, *J. Am. Chem. Soc.*, 1994, **116**, 7642.
- 14 M. S. Newman, R. S. Darlak and L. Tsai, *J. Am. Chem. Soc.*, 1967, **89**, 6191.
- 15 S. R. Wilson, Q. Lu, J. Cao, Y. Wu, C. J. Welch and D. I. Schuster, *Tetrahedron*, 1996, **52**, 5131.
- 16 M. Maggini, G. Scorrano, A. Bianco, C. Toniolo, R. P. Sijbesma, F. Wudl and M. Prato, *J. Chem. Soc., Chem. Commun.*, 1994, 305.
- 17 F. Novello, M. Prato, T. Da Ros, M. De Amici, A. Bianco, C. Toniolo and M. Maggini, *Chem. Commun.*, 1996, 903.

Received in Cambridge, UK, 8th October 1997; 7/07274K